

# What Is Allopurinol And Which are Allopurinol Side Effects

**W**hat Is Allopurinol And Which are Allopurinol Side effects in an effort to help people who are thinking to buy allopurinol to be as inform as They are able to before taking this product.



“ *Zyloprim is used in the treatment of many symptoms of gout, including serious attacks, tophi (collection of uric acid crystals in the cells, especially around joints), joint destruction, and uric acid stones. Gout is actually a form of arthritis characterized by increased blood levels of uric acid. Zyloprim works by decreasing uric acid production in the body, thus preventing crystals from forming. Zyloprim also manages the increased uric acid levels in the blood of individuals with certain cancers, for example leukemia.*

## Is Also Prescribed to Handle Some Types of Kidney Stones

Zyloprim it is not going to stop a gout strike that is already underway. However, when taken over a period of several months, Zyloprim it will reduce your symptoms. It's important to keep taking it regularly, even if it seems to have no immediate effect. Just how you should take Zyloprim Some low dosage and then he or she will keep upon increasing it every week or so until your reach the perfect dose for you. A normal starting dose is 100 MG Take Zyloprim exactly as approved by your doctor, he will start by recommending you one tablet per day, additionally you may want to take Zyloprim after a meal to reduce the risk of stomach irritation It can be not recommended that you do not get large doses of Vitamin C as it will increase the possibility of kidney stone formation While taking Zyloprim you should drink a lot of liquids--10 to 12 glasses (8 ounces each) per day-- unless otherwise prescribed by your doctor.

Order to help prevent attacks of gout, you should also avoid beer, wine, and also purine-rich foods such as anchovies, sardines, liver, kidneys, lentils, and also sweetbreads. If you have been taking colchicine and/or a great anti-inflammatory drug, such as Anaprox, Indocin, and others, to relieve your own gout, your doctor will probably want you to continue taking Zyloprim while the Zyloprim dosage is being adjusted. Later, when you have had no attacks of gout for many months, you may be able to stop taking other drugs.

You have been going for a drug that promotes the excretion of uric acid in the urine, such as probenecid (Benemid) or sulfinpyrazone (Anturane), to try to prevent attacks of gout, your physician will probably want to reduce or stop your dosage of Zyloprim while increasing your dosage of Zyloprim. If you miss a dose When you have forgotten to take a dose, take it as soon as you recall. If it is almost time for the next dose, skip the missed one and get back to your regular schedule. Do not take two doses immediately.

Zyloprim Side Effects Some medicines may cause side effects on some people but some may not, please check with your doctor if any of these side effects may persist or turn out to be irritating. Diarrhea; nausea. Find medical attention right away in the event that any of these SEVERE negative effects occur: Serious allergic reactions (rash; hives; itching; difficulty breathing; tightness in the chest; puffiness from the mouth, face, lip area, or tongue); blood in urine or painful urination; dark urine; fever, chills, or sore throat; irritation of the eyes; joint pain; lack of appetite; red, swollen, blistered, or peeling skin; stomach pain; unexplained weight loss; unusual bruising or bleeding; unusual muscle soreness or weakness; yellowing of the skin or eyes.



GoutUric AcidUricGout AttacksUric Acid LevelsGout DoctorJoint

## This is Not a Complete List of All Side Effects that May Occur

Crucial Zyloprim may cause drowsiness. This effect may be worse if you take it with alcohol or specific medicines. Use Zyloprim with caution. Do not drive or perform other possibly dangerous tasks until you know how you react to it. You may experience a great increase in gout attacks for many months after you begin taking Zyloprim. Your doctor may give you another medicine (colchicine) to help reduce these gout attacks.

- Contact your doctor if you go through frequent or severe gout attacks.
- It could take 2 to six months to be able to receive the full benefit of Zyloprim.
- Do NOT take more than the recommended dose or use for longer than recommended without checking with your doctor.
- Zyloprim is not a good choice for reducing an acute gout attack.
- Follow the diet and exercise program given to you by your health care provider.
- Lab tests, including uric acid levels and kidney and liver function, may be performed while you use Zyloprim.

## These Tests May be Used to Monitor Your Condition or Check for Side Effects

Make sure you keep all doctor and lab appointments. PREGNANCY and BREAST-FEEDING: If you become pregnant, contact your doctor. You will need to talk about the benefits and risks of using Zyloprim while you're pregnant. Zyloprim is found in breast milk. If you are or is going to be breast-feeding when you use Zyloprim, check with your doctor. Discuss any possible risks to be able to your baby.

Wrote the What is Allopurinol And Which are Allopurinol Side Effects in an effort to help people who are thinking to buy allopurinol being as inform as They could before taking this product.



### “ Domenic Golden

*Domenic is a head content marketing specialist at musclenstress.com, a collection of articles on health issues. In the past, Domenic worked as a post curator for a well-known health site. When he's not writing posts, Domenic enjoys drawing and rock climbing.*

**Disclaimer:** Content in this document is an advertisement, therefore it may be biased and should not be considered an objective or independent review. Owner of this document may be compensated when you purchase a product by clicking a link in this document. The views and opinions expressed in this document are purely of the author. Any product claim or other representation about a product or service should be verified with the manufacturer, provider or party in question. Any health information in this document is for educational purposes only and is not intended to replace the advice of your health care provider.